



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The two isosceles triangles CJE and IJK , having equal angles at E and K respectively, are similar. Hence we have $JE : CE :: JK : IK$, or, using (1), (2) and (3)

$$h\sqrt{4 - h^2} : (3h - h^3) :: h : (2h - h^3).$$

Hence

$$(4') \quad (2 - h^2)\sqrt{4 - h^2} = 3 - h^2.$$

Squaring, we get

$$(4 - 4h^2 + h^4)(4 - h^2) = 9 - 6h^2 + h^4,$$

and expanding,

$$16 - 20h^2 + 8h^4 - h^6 = 9 - 6h^2 + h^4.$$

Finally, transposing and simplifying, we obtain the author's *Heptagon Cubic*:

$$(5) \quad 7 - 14h^2 + 7h^4 - h^6 = 0.$$

Solving by Horner's method, we find $h^2 = .7530203962821 \dots = \frac{3}{4}$, approximately.

Remark. It will thus be seen that while there is introduced a new line IJ , we dispense with the consideration of the line OC , and with both the consideration and the computation of the author's lines SE , SJ , SK and SC . As a result, the equation (4') appears in a much simpler form than the author's equation (4).

The approximate construction of the heptagon may also be simplified as follows:

Let M , N and P be three consecutive vertices of an inscribed regular hexagon. Draw the chord MP and the radius ON , and let MP meet ON in R . Then MR is, approximately, the length h of the side of the regular inscribed heptagon. The reason is self-evident: approximately, $h = \frac{1}{2}\sqrt{3}$, and MP , as the side of a regular inscribed triangle, $= \sqrt{3}$, so that $MR = \frac{1}{2}\sqrt{3}$, and therefore $MR = h$, approximately.

A PROBLEM IN NUMBER THEORY.

By GEO. A. OSBORNE, Massachusetts Institute of Technology.

§ 1. When is the sum of the squares of two successive integers a perfect square? The following are examples:

$$3^2 + 4^2 = 5^2, \quad 20^2 + 21^2 = 29^2. \quad \text{The next is } 119^2 + 120^2 = 169^2.$$

The numbers 3, 20, 119, . . . are the terms of a series

$$0, 3, 20, 119, 696, \dots u_n, u_{n+1}, \quad (1)$$

where

$$u_{n+1} = 6u_n - u_{n-1} + 2. \quad (2)$$

This may be proved as follows:

From (2), which is the relation between any three successive terms of (1), we may derive the relation between any two successive terms as follows:

From (2)

$$\begin{aligned}u_{n+1} + u_{n-1} &= 6u_n + 2, \\u_{n+1}^2 - u_{n-1}^2 &= (6u_n + 2)(u_{n+1} - u_{n-1}), \\(u_{n+1} - 1)^2 - 6u_n u_{n+1} &= (u_{n-1} - 1)^2 - 6u_n u_{n-1},\end{aligned}$$

Adding to each member $(u_n - 1)^2$, we have

$$(u_{n+1} - 1)^2 + (u_n - 1)^2 - 6u_n u_{n+1} = (u_n - 1)^2 + (u_{n-1} - 1)^2 - 6u_{n-1} u_n, \quad (3)$$

which is of the form $f(n) = f(n-1)$.

Hence by induction, $f(n) = c$, a constant independent of n . By applying the first member of (3) to the terms 3 and 20, we find $c = 5$. Hence

$$(u_{n+1} - 1)^2 + (u_n - 1)^2 - 6u_n u_{n+1} = 5 \quad (4)$$

is the relation between any two successive terms of (1).

Solving (4) with respect to u_{n+1} , we have

$$u_{n+1} = 3u_n + 1 \pm 2\sqrt{2u_n^2 + 2u_n + 1}, \quad (5)$$

in which the lower sign is rejected since, otherwise, the right member would be less than u_n . It follows from (5) that

$$2u_n^2 + 2u_n + 1 = \text{a square},$$

that is,

$$u_n^2 + (u_n + 1)^2 = \text{a square},$$

one part of the result which was to be proved.

§ 2. From the terms of (1) we may write

$$\begin{aligned}0^2 + 1^2 &= 1^2, \\3^2 + 4^2 &= 5^2, \\20^2 + 21^2 &= 29^2, \\119^2 + 120^2 &= 169^2, \\696^2 + 697^2 &= 985^2, \\4059^2 + 4060^2 &= 5741^2, \\23660^2 + 23661^2 &= 33461^2, \\137903^2 + 137904^2 &= 195025^2, \\803760^2 + 803761^2 &= 1136689^2, \\\cdot &\quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \\\cdot &\quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot\end{aligned}$$

The second members are terms of a series,

$$1, 5, 29, \dots u_n, u_{n+1},$$

where

$$u_{n+1} = 6u_n - u_{n-1}.$$

§ 3. It remains to be shown that the terms of the series (1) are the *only* integers that satisfy the condition

$$N^2 + (N + 1)^2 = \text{a square.} \quad (6)$$

If from (4) we express u_n in terms of u_{n+1} , we have

$$u_n = 3u_{n+1} + 1 - 2\sqrt{2u_{n+1}^2 + 2u_{n+1} + 1}, \quad (7)$$

from which

$$u_{n-1} = 3u_n + 1 - 2\sqrt{2u_n^2 + 2u_n + 1}. \quad (8)$$

Consider the equation

$$y = 3x + 1 - 2\sqrt{2x^2 + 2x + 1}. \quad (9)$$

Then

$$\begin{aligned} 2y^2 + 2y + 1 &= 2(3x + 1 - 2\sqrt{2x^2 + 2x + 1})^2 + 2(3x + 1 - 2\sqrt{2x^2 + 2x + 1}) + 1 \\ &= (4x + 2 - 3\sqrt{2x^2 + 2x + 1})^2. \end{aligned}$$

Hence if

$$2x^2 + 2x + 1 = \text{a square,}$$

then also

$$2y^2 + 2y + 1 = \text{a square.}$$

That is, if x satisfies (6), so does y . Comparing (9) with (7) and (8), it appears that if $x = u_{n+1}$, $y = u_n$; and if $x = u_n$, $y = u_{n-1}$. And as y is an increasing function of x , since

$$\frac{dy}{dx} = 3 - \frac{4x + 2}{\sqrt{2x^2 + 2x + 1}} > 0,$$

it follows that if $u_n < x < u_{n+1}$, then $u_{n-1} < y < u_n$. That is, if there is an integer satisfying (6) between u_n and u_{n+1} , there is another such integer between u_{n-1} and u_n .

There is no integer satisfying (6) between the terms 3 and 20; hence there is none between 20 and 119, and consequently none between any two successive terms of the series (1).

§ 4. In the list of equations in § 2, it may be noticed that none of the successive integers end in 2, 5 or 8. As the final digits recur, it follows that

The sum of the squares of two successive integers, one of which ends in 2, 5 or 8, cannot be a perfect square.